

PCDA6 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56566**Specification**

PCDA6 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	O9UN73
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	102716

PCDA6 Polyclonal Antibody - Additional Information**Gene ID** 56142**Other Names**

Protocadherin alpha-6, PCDH-alpha-6, PCDHA6, CNRS2

Dilution

IHC-P ~ ~ N/A
IHC-F ~ ~ N/A
IF ~ ~ 1:50 ~ 200
E ~ ~ N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PCDA6 Polyclonal Antibody - Protein Information**Name** PCDHA6**Synonyms** CNRS2**Function**

Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.

Cellular Location


[Isoform 1]: Cell membrane; Single- pass type I membrane protein

PCDA6 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PCDA6 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PCDA6) Polyclonal Antibody, Unconjugated (bs-1711R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.