

TRMT12 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56576**Specification**

TRMT12 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q53H54
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50236

TRMT12 Polyclonal Antibody - Additional Information**Gene ID** 55039**Other Names**

tRNA wybutosine-synthesizing protein 2 homolog, tRNA-yW-synthesizing protein 2, 2.5.1.114, tRNA(Phe) (4-demethylwyosine(37)-C(7)) aminocarboxypropyltransferase, TRMT12, TRM12, TYW2

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

TRMT12 Polyclonal Antibody - Protein Information**Name** TRMT12**Synonyms** TRM12, TYW2**Function**


S-adenosyl-L-methionine-dependent transferase that acts as a component of the wybutosine biosynthesis pathway. Wybutosine is a hyper modified guanosine with a tricyclic base found at the 3'-position adjacent to the anticodon of eukaryotic phenylalanine tRNA. Catalyzes the transfer of the alpha-amino-alpha-carboxypropyl (acp) group from S- adenosyl-L-methionine to the C-7 position of 4-demethylwyosine (imG-14) to produce wybutosine-86.

TRMT12 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRMT12 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat ovary); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRMT12) Polyclonal Antibody, Unconjugated (bs-17144R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Sample: Ovary (Mouse) Lysate at 40 ug
Primary: Anti- TRMT12-1124 (bs-17144R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 50 kD
Observed band size: 50 kD