

IWS1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56598

Specification

IWS1 Polyclonal Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

IHC-P, IHC-F, IF, ICC <u>096ST2</u> Rat, Pig, Bovine Rabbit

Polyclonal 91955

IWS1 Polyclonal Antibody - Additional Information

Gene ID 55677

Other Names

Protein IWS1 homolog, IWS1-like protein, IWS1, IWS1L

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

IWS1 Polyclonal Antibody - Protein Information

Name IWS1

Synonyms IWS1L

Function

Transcription factor which plays a key role in defining the composition of the RNA polymerase II (RNAPII) elongation complex and in modulating the production of mature mRNA transcripts. Acts as an assembly factor to recruit various factors to the RNAPII elongation complex and is recruited to the complex via binding to the transcription elongation factor SUPT6H bound to the C-terminal domain (CTD) of the RNAPII subunit RPB1 (POLR2A). The SUPT6H:IWS1:CTD complex recruits mRNA export factors (ALYREF/THOC4, EXOSC10) as well as histone modifying enzymes (such as SETD2) to ensure proper mRNA splicing, efficient mRNA export and elongation-coupled H3K36 methylation, a signature chromatin mark of active transcription.

Cellular Location

 $Nucleus \ \{ ECO: 0000255 | PROSITE-ProRule: PRU00649, \ ECO: 0000269 | PubMed: 17184735, \ ECO: 0000269 | PubMed: 17234882 \}$



Tel: 858.875.1900 Fax: 858.875.1999



IWS1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

IWS1 Polyclonal Antibody - Images