

SAFB Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56629**Specification**

SAFB Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q14151
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	107473

SAFB Polyclonal Antibody - Additional Information**Gene ID** 9667**Other Names**

Scaffold attachment factor B2, SAF-B2, SAFB2, KIAA0138

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SAFB Polyclonal Antibody - Protein Information**Name** SAFB2**Synonyms** KIAA0138**Function**

Binds to scaffold/matrix attachment region (S/MAR) DNA. Can function as an estrogen receptor corepressor and can also inhibit cell proliferation.

Cellular Location

Cytoplasm. Nucleus.

Tissue Location


Expressed at high levels in the CNS and at low levels in the liver. Expressed in a wide number of breast cancer cell lines

SAFB Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SAFB Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SAFB) Polyclonal Antibody, Unconjugated (bs-17259R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SAFB) Polyclonal Antibody, Unconjugated (bs-17259R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.