

**SFRS2IP Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56651****Specification**

---

**SFRS2IP Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">Q99590</a>
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	164652

**SFRS2IP Polyclonal Antibody - Additional Information****Gene ID** 9169**Other Names**

Protein SCAF11, CTD-associated SR protein 11, Renal carcinoma antigen NY-REN-40, SC35-interacting protein 1, SR-related and CTD-associated factor 11, SRSF2-interacting protein, Serine/arginine-rich splicing factor 2-interacting protein, Splicing factor, arginine/serine-rich 2-interacting protein, Splicing regulatory protein 129, SRp129, SCAF11, CASP11, SFRS2IP, SIP1, SRSF2IP

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**SFRS2IP Polyclonal Antibody - Protein Information****Name** SCAF11**Synonyms** CASP11, SFRS2IP, SIP1, SRSF2IP**Function**

Plays a role in pre-mRNA alternative splicing by regulating spliceosome assembly.

**Cellular Location**

Nucleus.

**Tissue Location**


Widely expressed..

## SFRS2IP Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## SFRS2IP Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat lymphoid tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SFRS2IP) Polyclonal Antibody, Unconjugated (bs-17298R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse testis tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SFRS2IP) Polyclonal Antibody, Unconjugated (bs-17298R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.