

**SFRS5 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56652****Specification**

---

**SFRS5 Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">Q13243</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31264

**SFRS5 Polyclonal Antibody - Additional Information****Gene ID** 6430**Other Names**

Serine/arginine-rich splicing factor 5, Delayed-early protein HRS, Pre-mRNA-splicing factor SRP40, Splicing factor, arginine/serine-rich 5, SRSF5, HRS, SFRS5, SRP40

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**SFRS5 Polyclonal Antibody - Protein Information****Name** SRSF5**Synonyms** HRS, SFRS5, SRP40**Function**

Plays a role in constitutive splicing and can modulate the selection of alternative splice sites.

**Cellular Location**

Nucleus.


**SFRS5 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **SFRS5 Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SFRS5) Polyclonal Antibody, Unconjugated (bs-17301R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.