

### hnRNP A0 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56662

#### Specification

## hnRNP A0 Polyclonal Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW Physical State Immunogen Epitope Specificity Isotype <b>Purity</b> affinity purified by Protein A	WB, IHC-P, IHC-F, IF, ICC, E <u>O13151</u> Rat, Pig, Cynomolgus, Dog, Bovine Rabbit Polyclonal 31 KDa Liquid KLH conjugated synthetic peptide derived from human hnRNP A0 1-100/305 IgG
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION SIMILARITY	Nucleus. Component of ribonucleosomes. Contains 2 RRM (RNA recognition motif) domains.
Post-translational modifications	Arg-291 is dimethylated, probably to asymmetric dimethylarginine.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

#### **Background Descriptions**

Heterogeneous nuclear ribonucleoproteins (hnRNPs) constitute a set of polypeptides that contribute to mRNA transcription and pre-mRNA processing as well as mature mRNA transport to the cytoplasm and translation. They also bind heterogeneous nuclear RNA (hnRNA), which are the transcripts produced by RNA polymerase II. There are approximately 20 known hnRNP proteins and their complexes are the major constituents of the spliceosome. The majority of hnRNP protein components are localized to the nucleus; however, some shuttle between the nucleus and the cytoplasm. hnRNP I, also designated polypyrimidine tract-binding protein (PTB), and its homolog hnRNP L, bind to the 3' end of introns to modulate alternative splicing mechanisms of pre-mRNAs in normal cells and the translation of sev-eral viruses, including hepatitis C virus (HCV). The human hnRNP I gene maps to chromosome 19p13.3 and encodes a protein that is localized in the nucleoplasm. hnRNP L, like hnRNP I, is also localized in the nucleoplasm.

#### hnRNP A0 Polyclonal Antibody - Additional Information

Gene ID 10949

**Other Names** Heterogeneous nuclear ribonucleoprotein A0, hnRNP A0, HNRNPA0, HNRPA0



Dilution

<span class ="dilution\_WB">WB~~1:1000</span><br \><span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \><span class ="dilution\_IHC-F">IHC-F~~N/A</span><br \><span class ="dilution\_IF">IF~~1:50~200</span><br \><span class ="dilution\_ICC">ICC~~N/A</span><br \><span class = "dilution\_ICC">ICC~~N/A</span><br \><span class = "dilution\_ICC">ICC~~N/A</sp

Format 0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

# hnRNP A0 Polyclonal Antibody - Protein Information

Name HNRNPA0

Synonyms HNRPA0

Function

mRNA-binding component of ribonucleosomes. Specifically binds AU-rich element (ARE)-containing mRNAs. Involved in post- transcriptional regulation of cytokines mRNAs.

**Cellular Location** 

Nucleus. Note=Component of ribonucleosomes.

#### hnRNP A0 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

hnRNP A0 Polyclonal Antibody - Images





Sample:

Raw264.7(Mouse) Cell Lysate at 30 ug Primary: Anti-hnRNP A0 (bs-17339R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 31 kD Observed band size: 31 kD