

hnRNP H3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56663**Specification**

hnRNP H3 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	P31942
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36926

hnRNP H3 Polyclonal Antibody - Additional Information**Gene ID** 3189**Other Names**

Heterogeneous nuclear ribonucleoprotein H3, hnRNP H3, Heterogeneous nuclear ribonucleoprotein 2H9, hnRNP 2H9, HNRNPH3, HNRPH3

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

hnRNP H3 Polyclonal Antibody - Protein Information**Name** HNRNPH3**Synonyms** HNRPH3**Function**

Involved in the splicing process and participates in early heat shock-induced splicing arrest. Due to their great structural variations the different isoforms may possess different functions in the splicing reaction.

Cellular Location


Nucleus.

hnRNP H3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

hnRNP H3 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat uterus); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (hnRNP H3) Polyclonal Antibody, Unconjugated (bs-17345R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.