

**HOOK3 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56668****Specification**

---

**HOOK3 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q86VS8</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	83126

**HOOK3 Polyclonal Antibody - Additional Information****Gene ID** 84376**Other Names**

Protein Hook homolog 3, h-hook3, hHK3, HOOK3

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**HOOK3 Polyclonal Antibody - Protein Information****Name** HOOK3 ([HGNC:23576](#))**Function**

Acts as an adapter protein linking the dynein motor complex to various cargos and converts dynein from a non-processive to a highly processive motor in the presence of dynactin. Facilitates the interaction between dynein and dynactin and activates dynein processivity (the ability to move along a microtubule for a long distance without falling off the track). Predominantly recruits 2 dyneins, which increases both the force and speed of the microtubule motor (PubMed:<a href="http://www.uniprot.org/citations/25035494" target="\_blank">25035494</a>, PubMed:<a href="http://www.uniprot.org/citations/33734450" target="\_blank">33734450</a>). Component of the FTS/Hook/FHIP complex (FHF complex). The FHF complex may function to promote vesicle trafficking and/or fusion via the homotypic vesicular protein sorting complex (the HOPS complex). May regulate clearance of endocytosed receptors such as MSR1. Participates in defining the architecture and localization of the Golgi complex. FHF complex promotes the distribution of AP-4 complex to the perinuclear area of the cell (PubMed:<a href="http://www.uniprot.org/citations/32073997" target="\_blank">32073997</a>).

**Cellular Location**

Cytoplasm, cytoskeleton. Golgi apparatus. Note=Enriched at the cis-face of the Golgi complex.


Localizes to microtubule asters in prophase (PubMed:11238449). Localizes to the manchette in elongating spermatids (By similarity). {ECO:0000250|UniProtKB:Q8BUK6, ECO:0000269|PubMed:11238449}

### HOOK3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### HOOK3 Polyclonal Antibody - Images



**Sample:**

Small intestine(Mouse) Cell Lysate at 40 ug

Primary: Anti-HOOK3(bs-17355R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 83 kD

Observed band size: 83 kD