

SPIRE1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56791**Specification****SPIRE1 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q08AE8
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85544

SPIRE1 Polyclonal Antibody - Additional Information**Gene ID** 56907**Other Names**

Protein spire homolog 1, Spir-1, SPIRE1 {ECO:0000312|EMBL:AAI25207.1}, KIAA1135, SPIR1

Dilution

IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SPIRE1 Polyclonal Antibody - Protein Information**Name** SPIRE1 {ECO:0000312|EMBL:AAI25207.1}**Synonyms** KIAA1135, SPIR1**Function**

Acts as an actin nucleation factor, remains associated with the slow-growing pointed end of the new filament (PubMed:11747823, PubMed:21620703). Involved in intracellular vesicle transport along actin fibers, providing a novel link between actin cytoskeleton dynamics and intracellular transport (PubMed:11747823). Required for asymmetric spindle positioning and asymmetric cell division during meiosis (PubMed:21620703). Required for normal formation of the cleavage furrow and for polar body extrusion during female germ cell meiosis (PubMed:21620703). Also acts in the nucleus: together with FMN2, promotes

assembly of nuclear actin filaments in response to DNA damage in order to facilitate movement of chromatin and repair factors after DNA damage (PubMed:26287480). In addition, promotes innate immune signaling downstream of dsRNA sensing (PubMed:35148361). Mechanistically, contributes to IRF3 phosphorylation and activation downstream of MAVS and upstream of TBK1 (PubMed:35148361).

Cellular Location

Cytoplasm, cytoskeleton. Cytoplasm, perinuclear region. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q52KF3}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q52KF3}; Cytoplasmic side {ECO:0000250|UniProtKB:Q52KF3}. Note=Detected at the cleavage furrow during asymmetric oocyte division and polar body extrusion (By similarity). Punctate spots in perinuclear region and cytoplasm, colocalized with Rab11 (By similarity). {ECO:0000250|UniProtKB:Q52KF3}

SPIRE1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SPIRE1 Polyclonal Antibody - Images