



## **MRP6 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) **Catalog # AP56823** 

### **Specification**

# MRP6 Polyclonal Antibody - Product Information

Application **Primary Accession** 

Reactivity Host Clonality Calculated MW **Physical State** 

Immunogen

**Epitope Specificity** 

Isotype **Purity** 

affinity purified by Protein A

Buffer

SUBCELLULAR LOCATION

**SIMILARITY** 

**DISEASE** 

Important Note

**Background Descriptions** 

IHC-P, IHC-F, IF, ICC, E

095255

Rat, Pig, Bovine

**Rabbit Polyclonal** 165 KDa Liquid

KLH conjugated synthetic peptide derived

from human MRP6 1351-1503/1503

laG

0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

Membrane. Localized to the basolateral

membrane.

**Belongs to the ABC transporter** superfamily. ABCC family. Conjugate transporter (TC 3.A.1.208) subfamily. Contains 2 ABC transmembrane type-1 domains. Contains 2 ABC transporter

domains.

Defects in ABCC6 are the cause of pseudoxanthoma elasticum (PXE) [MIM:264800]. PXE is a disorder

characterized by calcification of elastic fibers in skin, arteries and retina that results in dermal lesions with associated laxity and loss of elasticity, arterial insufficiency and retinal hemorrhages leading to macular degeneration. PXE is caused in the overwhelming majority of cases by homozygous or compound

heterozygous mutations in the ABCC6 gene (autosomal recessive PXE). Individuals carrying heterozygous mutations express limited manifestations of the

pseudoxanthoma elasticum phenotype

(autosomal dominant PXE).

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). The encoded protein, a member of the MRP subfamily, is involved in multi-drug resistance. Mutations in this gene cause pseudoxanthoma elasticum. Alternatively spliced transcript variants that encode different proteins have been described for this gene. [provided by RefSeq, Jul 2008]

## MRP6 Polyclonal Antibody - Additional Information

Gene ID 368

### **Other Names**

ATP-binding cassette sub-family C member 6, 7.6.2.-, 7.6.2.3, Anthracycline resistance-associated protein, Multi-specific organic anion transporter E, MOAT-E, Multidrug resistance-associated protein 6, ABCC6, ARA, MRP6

# Target/Specificity

Expressed in kidney and liver. Very low expression in other tissues.

### **Dilution**

```
<span class ="dilution_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution_IF">IF~~1:50~200</span><br \> <span class ="dilution_ICC">ICC~~N/A</span><br \> <span class ="dilution_E">E~~N/A</span>
```

## **Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## **MRP6 Polyclonal Antibody - Protein Information**

Name ABCC6

Synonyms ARA, MRP6

#### **Function**

[Isoform 1]: ATP-dependent transporter of the ATP-binding cassette (ABC) family that actively extrudes physiological compounds, and xenobiotics from cells. Mediates ATP-dependent transport of glutathione conjugates such as leukotriene-c4 (LTC4) and N- ethylmaleimide S-glutathione (NEM-GS) (in vitro), and an anionic cyclopentapeptide endothelin antagonist, BQ-123 (PubMed:<a href="http://www.uniprot.org/citations/11880368" target="\_blank">11880368</a>, PubMed:<a href="http://www.uniprot.org/citations/12414644" target="\_blank">12414644</a>). May contribute to regulate the transport of organic compounds in testes across the blood-testis-barrier (Probable). Does not appear to actively transport drugs outside the cell. Confers low levels of cellular resistance to etoposide, teniposide, anthracyclines and cisplatin (PubMed:<a href="http://www.uniprot.org/citations/12414644" target="\_blank">12414644</a>).

### **Cellular Location**

Basal cell membrane; Multi-pass membrane protein. Note=Localized to the basal membrane of Sertoli cells [Isoform 2]: Endoplasmic reticulum membrane; Single-pass membrane protein

## **Tissue Location**



Expressed in kidney and liver. Very low expression in other tissues. In testis, localized to peritubular myoid cells, Leydig cells, along the basal membrane of Sertoli cells and moderately in the adluminal compartment of the seminiferous tubules (PubMed:35307651).

# **MRP6 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

**MRP6 Polyclonal Antibody - Images**