

MRPL47 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56837**Specification**

MRPL47 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	O9HD33
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29450

MRPL47 Polyclonal Antibody - Additional Information**Gene ID** 57129**Other Names**

39S ribosomal protein L47, mitochondrial, L47mt, MRP-L47, Mitochondrial large ribosomal subunit protein uL29m, Nasopharyngeal carcinoma metastasis-related protein 1, MRPL47, NCM1

Dilution

IHC-P ~ ~ N/A
IHC-F ~ ~ N/A
IF ~ ~ 1:50 ~ 200
ICC ~ ~ N/A
E ~ ~ N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

MRPL47 Polyclonal Antibody - Protein Information**Name** MRPL47**Synonyms** NCM1**Cellular Location**


Mitochondrion

MRPL47 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MRPL47 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MRPL47) Polyclonal Antibody, Unconjugated (bs-17800R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.