

**CD20L4/MS4A7 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56862****Specification**

---

**CD20L4/MS4A7 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">O9GZW8</a>
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26131

**CD20L4/MS4A7 Polyclonal Antibody - Additional Information****Gene ID** 58475**Other Names**

Membrane-spanning 4-domains subfamily A member 7, CD20 antigen-like 4, CD20/FC-epsilon-RI-beta family member 4, Four-span transmembrane protein 2, MS4A7, 4SPAN2, CD20L4, CFFM4

**Dilution**

IHC-P ~ ~ N/A  
IHC-F ~ ~ N/A  
IF ~ ~ 1:50 ~ 200  
ICC ~ ~ N/A  
E ~ ~ N/A

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glycerol

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**CD20L4/MS4A7 Polyclonal Antibody - Protein Information****Name** MS4A7**Synonyms** 4SPAN2, CD20L4, CFFM4**Function**

May be involved in signal transduction as a component of a multimeric receptor complex.

**Cellular Location**

Membrane; Multi-pass membrane protein.

**Tissue Location**

Ubiquitous expression in normal tissues. Expression is more elevated in adult liver, lung, spleen,

and heart than in their fetal counterparts, and is higher in normal tissues than in the cancerous tissue or cell lines. Low levels of expression were detected in the promonocytic stage, whereas high levels of expression were detected in mature monocytes

### **CD20L4/MS4A7 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **CD20L4/MS4A7 Polyclonal Antibody - Images**