

## CXCL6/GCP2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56906

#### **Specification**

## CXCL6/GCP2 Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, E

Primary Accession
Reactivity
Bovine
Host
Clonality
Calculated MW
Physical State
P50228
Rabbit
Rabbit
Polyclonal
12 KDa
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human CXCL6

Epitope Specificity 51-114/114

Isotype IgG

**Purity**affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Secreted.

SIMILARITY Belongs to the intercrine alpha (chemokine

CxC) family.

Important Note This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

#### **Background Descriptions**

Neutrophil recruitment to inflammatory sites is mediated by two related receptors: CXCR1 and CXCR2. Both receptors share two ligands, interleukin-8 (CXCL8) and GCP2 (CXCL6). GCP2 belongs to the intercrine alpha (chemokine CxC) family.

## CXCL6/GCP2 Polyclonal Antibody - Additional Information

#### Gene ID 20311

# **Other Names**

C-X-C motif chemokine 5, Cytokine LIX, Small-inducible cytokine B5, GCP-2(1-78), GCP-2(9-78), Cxcl5, Scyb5

## **Dilution**

<span class ="dilution WB">WB~~1:1000</span><br \><span class</pre>

="dilution IHC-P">IHC-P~~N/A</span><br \><span class

="dilution IHC-F">IHC-F~~N/A</span><br \><span class

="dilution IF">IF $\sim$ 1:50 $\sim$ 200</span><br\><span class ="dilution E">E $\sim$ N/A</span>

#### **Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce



## **Storage**

Store at -20  $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.

# CXCL6/GCP2 Polyclonal Antibody - Protein Information

Name Cxcl5

Synonyms Scyb5

#### **Function**

May participate in the recruitment of inflammatory cells by injured or infected tissue. GCP-2(1-78) and, more potent, GCP-2(9-78) attract neutrophils and are involved in neutrophil activation.

**Cellular Location** 

Secreted.

# CXCL6/GCP2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# CXCL6/GCP2 Polyclonal Antibody - Images