

FBXO34 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5692b**Specification**

FBXO34 Antibody (C-term) - Product Information

Application	FC, IHC-P, WB,E
Primary Accession	Q9NWN3
Other Accession	Q80XI1 , NP_060413.2
Reactivity	Human, Hamster
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	78711
Antigen Region	600-629

FBXO34 Antibody (C-term) - Additional Information**Gene ID** 55030**Other Names**

F-box only protein 34, FBXO34, FBX34

Target/Specificity

This FBXO34 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 600-629 amino acids from the C-terminal region of human FBXO34.

Dilution

FC~~1:10~50

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FBXO34 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FBXO34 Antibody (C-term) - Protein Information

Name FBXO34

Synonyms FBX34


Function Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex.

FBXO34 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FBXO34 Antibody (C-term) - Images




FBXO34 Antibody (C-term) (Cat. #AP5692b) western blot analysis in CHO cell line lysates (35ug/lane). This demonstrates the FBXO34 antibody detected the FBXO34 protein (arrow).



FBXO34 Antibody (C-term) (Cat. #AP5692b) immunohistochemistry analysis in formalin fixed and paraffin embedded human colon carcinoma followed by peroxidase conjugation of the secondary

antibody and DAB staining. This data demonstrates the use of the FBXO34 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



FBXO34 Antibody (C-term) (Cat. #AP5692b) flow cytometric analysis of WiDr cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

FBXO34 Antibody (C-term) - Background

Members of the F-box protein family, such as FBXO34, are characterized by an approximately 40-amino acid F-box motif. SCF complexes, formed by SKP1 (MIM 601434), cullin (see CUL1; MIM 603134), and F-box proteins, act as protein-ubiquitin ligases. F-box proteins interact with SKP1 through the F box, and they interact with ubiquitination targets through other protein interaction domains (Jin et al., 2004 [PubMed 15520277]). [supplied by OMIM].

FBXO34 Antibody (C-term) - References

Barrios-Rodiles, M., et al. Science 307(5715):1621-1625(2005)
Jin, J., et al. Genes Dev. 18(21):2573-2580(2004)