

HspBP1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56944**Specification****HspBP1 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	O9NZL4
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human HspBP1
Epitope Specificity	51-150/362
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 4 ARM repeats.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Hsp70-interacting protein (HspBP1) belongs to a family of eukaryotic proteins identified as nucleotide exchange factors for Hsp70, which exhibit varying degrees of compartment and species specificity. HspBP1 interferes with the CHIP-induced degradation of immature forms of the cystic fibrosis transmembrane conductance regulator (CFTR) and stimulates CFTR maturation. HspBP1 binds to Hsp70, inhibits its activity and promotes dissociation of nucleotides from the Hsp70 ATPase domain. It is a protein mainly expressed in heart and skeletal muscle.

HspBP1 Polyclonal Antibody - Additional Information**Gene ID** 23640**Other Names**

Hsp70-binding protein 1, HspBP1, Heat shock protein-binding protein 1, Hsp70-binding protein 2, HspBP2, Hsp70-interacting protein 1, Hsp70-interacting protein 2, HSPBP1 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=24989) target="_blank">HGNC:24989), HSPBP

Target/Specificity

Ubiquitous.

Dilution

IHC-P~~N/A<br \><span class

=IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

HspBP1 Polyclonal Antibody - Protein Information

Name HSPBP1 ([HGNC:24989](#))

Synonyms HSPBP

Function

Inhibits HSPA1A chaperone activity by changing the conformation of the ATP-binding domain of HSPA1A and interfering with ATP binding. Interferes with ubiquitination mediated by STUB1 and inhibits chaperone-assisted degradation of immature CFTR.


Tissue Location

Ubiquitous..

HspBP1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HspBP1 Polyclonal Antibody - Images

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HspBP1) Polyclonal Antibody, Unconjugated (bs-18085R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.