

RELT/TNFRSF19L Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56964**Specification**

RELT/TNFRSF19L Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	O969Z4
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46092

RELT/TNFRSF19L Polyclonal Antibody - Additional Information**Gene ID** 84957**Other Names**

Tumor necrosis factor receptor superfamily member 19L, Receptor expressed in lymphoid tissues, RELT, TNFRSF19L

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

RELT/TNFRSF19L Polyclonal Antibody - Protein Information**Name** RELT**Synonyms** TNFRSF19L**Function**

May play a role in apoptosis (PubMed:28688764, PubMed:19969290). Induces activation of MAPK14/p38 and MAPK8/JNK MAPK cascades, when overexpressed (PubMed:16530727). Involved in dental enamel formation (PubMed:30506946).

Cellular Location

Cell membrane; Single-pass type I membrane protein Cytoplasm. Cytoplasm, perinuclear region

Tissue Location

Spleen, lymph node, brain, breast and peripheral blood leukocytes (at protein level) (PubMed:28688764). Expressed highly in bone marrow and fetal liver. Very low levels in skeletal

muscle, testis and colon. Not detected in kidney and pancreas

RELT/TNFRSF19L Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RELT/TNFRSF19L Polyclonal Antibody - Images