

MLR2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP56986**Specification**

MLR2 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	O96JN0
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47007

MLR2 Polyclonal Antibody - Additional Information**Gene ID** 84458**Other Names**

Ligand-dependent corepressor, LCoR, Mblk1-related protein 2, LCOR (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=29503)
HGNC:29503

Dilution

IHC-P ~ N/A
IHC-F ~ N/A
IF ~ 1:50 ~ 200
ICC ~ N/A
E ~ N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

MLR2 Polyclonal Antibody - Protein Information**Name** LCOR ([HGNC:29503](#))**Function**

May act as transcription activator that binds DNA elements with the sequence 5'-CCCTATCGATCGATCTCTACCT-3' (By similarity). Repressor of ligand-dependent transcription activation by target nuclear receptors. Repressor of ligand-dependent transcription activation by ESR1, ESR2, NR3C1, PGR, RARA, RARB, RARG, RXRA and VDR.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00320, ECO:0000269|PubMed:12535528}

Tissue Location

Ubiquitous..

MLR2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MLR2 Polyclonal Antibody - Images