

**LENG8 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56997****Specification**

---

**LENG8 Polyclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | IHC-P                  |
| Primary Accession | <a href="#">Q96PV6</a> |
| Reactivity        | Rat, Pig, Bovine       |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Calculated MW     | 88157                  |

**LENG8 Polyclonal Antibody - Additional Information****Gene ID** 114823**Other Names**

Leukocyte receptor cluster member 8, LENG8, KIAA1932

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**


Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**LENG8 Polyclonal Antibody - Protein Information****Name** LENG8**Synonyms** KIAA1932**LENG8 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**LENG8 Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (Rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LENG8) Polyclonal Antibody, Unconjugated (bs-18220R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.