

**LINS Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP57020****Specification**

---

**LINS Polyclonal Antibody - Product Information**

Application	WB, IHC-P
Primary Accession	<a href="#">Q8NG48</a>
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85857

**LINS Polyclonal Antibody - Additional Information****Gene ID** 55180**Other Names**

Protein Lines homolog 1 {ECO:0000312|HGNC:HGNC:30922}, Wnt-signaling molecule Lines homolog 1, LINS1 ([HGNC:30922](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=30922))

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**LINS Polyclonal Antibody - Protein Information****Name** LINS1 ([HGNC:30922](#))**Tissue Location**

Expressed in adult testis, prostate, prostate, spleen, thymus, skeletal muscle, fetal kidney and brain


**LINS Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### LINS Polyclonal Antibody - Images



**Sample:**


U87MG(Human) Cell Lysate at 40 ug

Primary: Anti-LINS (bs-18286R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 86 kD

Observed band size: 86 kD



**Sample:**


U251(Human) Cell Lysate at 40 ug

Primary: Anti-LINS (bs-18286R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 86 kD

Observed band size: 86 kD



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LINS) Polyclonal Antibody, Unconjugated (bs-18286R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.