

## LMBR1L Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57031

## **Specification**

## LMBR1L Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC

Primary Accession <u>Q6UX01</u>

Reactivity Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 55 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

from human LMBR1L

Epitope Specificity 1-100/489

Isotype IgG
Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cell Membrane; Multi-pass membrane

protein.

SIMILARITY Belongs to the LIMR family.

Important Note

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

**Background Descriptions** 

LMBR1L is a probable LCN1 receptor and may mediate LCN1 endocytosis.

## LMBR1L Polyclonal Antibody - Additional Information

#### **Gene ID 55716**

#### **Other Names**

Protein LMBR1L, Limb region 1 protein homolog-like, Lipocalin-1-interacting membrane receptor, LIMR, LMBR1L, KIAA1174, LIMR

#### **Dilution**

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \><span class</pre>

="dilution\_IHC-F">IHC-F~~N/A</span><br \><span class

="dilution IF">IF~~1:50~200</span><br \><span class ="dilution ICC">ICC~~N/A</span>

#### **Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

## **Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH



7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

# LMBR1L Polyclonal Antibody - Protein Information

Name LMBR1L

Synonyms KIAA1174, LIMR

#### **Function**

Plays an essential role in lymphocyte development by negatively regulating the canonical Wnt signaling pathway (By similarity). In association with UBAC2 and E3 ubiquitin-protein ligase AMFR, promotes the ubiquitin-mediated degradation of CTNNB1 and Wnt receptors FZD6 and LRP6 (By similarity). LMBR1L stabilizes the beta- catenin destruction complex that is required for regulating CTNNB1 levels (By similarity). Acts as a LCN1 receptor and can mediate its endocytosis (PubMed:<a href="http://www.uniprot.org/citations/11287427" target="\_blank">11287427</a>, PubMed:<a href="http://www.uniprot.org/citations/12591932" target="\_blank">12591932</a>, PubMed:<a href="http://www.uniprot.org/citations/23964685" target="\_blank">23964685</a>).

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein

#### **Tissue Location**

Expressed in testis, pituitary gland, adrenal gland, trachea, placenta, thymus, cerebellum, stomach, mammary gland, spinal cord. A weaker expression is detected in colon, pancreas, and prostate.

## **LMBR1L Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## LMBR1L Polyclonal Antibody - Images