

LTV1/C6orf93 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57085**Specification**

LTV1/C6orf93 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC
Primary Accession	Q96GA3
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54855

LTV1/C6orf93 Polyclonal Antibody - Additional Information**Gene ID** 84946**Other Names**

Protein LTV1 homolog, LTV1, C6orf93

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

LTV1/C6orf93 Polyclonal Antibody - Protein Information**Name** LTV1**Synonyms** C6orf93**Function**

Essential for ribosome biogenesis.

Cellular Location

Cytoplasm. Note=The protein is expressed in a diffuse cytoplasmic pattern with granular perinuclear accentuation within the basal keratinocytes

Tissue Location


Expressed in the epidermis.

LTV1/C6orf93 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LTV1/C6orf93 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LTV1) Polyclonal Antibody, Unconjugated (bs-18442R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse placenta); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LTV1) Polyclonal Antibody, Unconjugated (bs-18442R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.