

**LUZP1 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP57086****Specification****LUZP1 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	<a href="#">Q86V48</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	120275

**LUZP1 Polyclonal Antibody - Additional Information****Gene ID 7798****Other Names**

Leucine zipper protein 1, LUZP1

**Dilution**

<span class = "dilution\_IHC-P">IHC-P~~N/A</span><br /><span class = "dilution\_IHC-F">IHC-F~~N/A</span><br /><span class = "dilution\_IF">IF~~1:50~200</span><br /><span class = "dilution\_ICC">ICC~~N/A</span>

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**LUZP1 Polyclonal Antibody - Protein Information****Name LUZP1****Function**

F-actin cross-linking protein (PubMed:<a href="http://www.uniprot.org/citations/30990684" target="\_blank">30990684</a>). Stabilizes actin and acts as a negative regulator of primary cilium formation (PubMed:<a href="http://www.uniprot.org/citations/32496561" target="\_blank">32496561</a>). Positively regulates the phosphorylation of both myosin II and protein phosphatase 1 regulatory subunit PPP1R12A/MYPT1 and promotes the assembly of myosin II stacks within actin stress fibers (PubMed:<a href="http://www.uniprot.org/citations/38832964" target="\_blank">38832964</a>). Inhibits the phosphorylation of myosin light chain MYL9 by DAPK3 and suppresses the constriction velocity of the contractile ring during cytokinesis (PubMed:<a href="http://www.uniprot.org/citations/38009294" target="\_blank">38009294</a>). Binds to microtubules and promotes epithelial cell apical constriction by up-regulating levels of diphosphorylated myosin light chain (MLC) through microtubule-dependent inhibition of MLC dephosphorylation by myosin phosphatase (By similarity). Involved in regulation of cell migration, nuclear size and centriole number, probably through regulation of the actin cytoskeleton (By similarity). Component of the CERF-1 and CERF-5 chromatin remodeling complexes in embryonic

stem cells where it acts to stabilize the complexes (By similarity). Plays a role in embryonic brain and cardiovascular development (By similarity).

### **Cellular Location**

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, cilium basal body. Midbody. Chromosome, centromere. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, stress fiber. Nucleus {ECO:0000250|UniProtKB:Q9ESV1} Cell projection, dendrite {ECO:0000250|UniProtKB:Q9ESV1}. Perikaryon {ECO:0000250|UniProtKB:Q9ESV1}. Cell junction, tight junction {ECO:0000250|UniProtKB:Q8R4U7}. Note=Localizes to the proximal end of basal bodies (PubMed:32496561). During mitosis, localizes at the inner centromere in metaphase, at the central spindle in anaphase, and at the midbody in telophase (PubMed:38009294). Central spindle localization requires KIF20A while centromere localization requires the kinase activity of the chromosomal passenger complex (PubMed:38009294)

### **LUZP1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **LUZP1 Polyclonal Antibody - Images**