

ZC3H12C Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57094**Specification**

ZC3H12C Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9C0D7
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	99340

ZC3H12C Polyclonal Antibody - Additional Information**Gene ID** 85463**Other Names**

Probable ribonuclease ZC3H12C, 3.1.-., MCP-induced protein 3, Zinc finger CCCH domain-containing protein 12C, ZC3H12C, KIAA1726, MCPIP3

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ZC3H12C Polyclonal Antibody - Protein Information**Name** ZC3H12C**Synonyms** KIAA1726, MCPIP3**Function**

May function as RNase and regulate the levels of target RNA species.


ZC3H12C Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZC3H12C Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZC3H12C) Polyclonal Antibody, Unconjugated (bs-18460R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.