

ZNF622 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP57123

Specification

ZNF622 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q969S3
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ZNF622
Epitope Specificity	1-100/477
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Nucleus.
SIMILARITY	Contains 2 U1-type zinc fingers.
Post-translational modifications	Phosphorylated by MELK. The phosphorylation may redirect the protein to the nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

ZNF622 (Zinc Finger Protein 622) is a Protein Coding gene. Diseases associated with ZNF622 include Autosomal Dominant Non-Syndromic Intellectual Disability 44 and Autoinflammation, Panniculitis, And Dermatitis Syndrome. Gene Ontology (GO) annotations related to this gene include nucleic acid binding.

ZNF622 Polyclonal Antibody - Additional Information

Gene ID 90441

Other Names

Zinc finger protein 622, Zinc finger-like protein 9, ZNF622, ZPR9

Target/Specificity

Expressed in lung, kidney, spleen, liver and brain with lowest expression in kidney.

Dilution

IHC-P ~ ~ N/A
IHC-F ~ ~ N/A

=IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ZNF622 Polyclonal Antibody - Protein Information

Name ZNF622 {ECO:0000303|PubMed:32669547, ECO:0000312|HGNC:HGNC:30958}

Function

Pre-60S-associated cytoplasmic factor involved in the cytoplasmic maturation of the 60S subunit.

Cellular Location

Cytoplasm. Nucleus

Tissue Location


Expressed in lung, kidney, spleen, liver and brain with lowest expression in kidney.

ZNF622 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF622 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF622) Polyclonal Antibody, Unconjugated (bs-18514R) at 1:200 overnight at 4°C, followed by

operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.