

ZFAT/ZNF406 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57128

Specification

ZFAT/ZNF406 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC

Primary Accession <u>O9P243</u>

Reactivity Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 139034

ZFAT/ZNF406 Polyclonal Antibody - Additional Information

Gene ID 57623

Other Names

Zinc finger protein ZFAT, Zinc finger gene in AITD susceptibility region, Zinc finger protein 406, ZFAT, KIAA1485, ZFAT1, ZNF406

Dilution

IHC-P~~N/A<br \> IHC-F~~N/A<br \> IF~~1:50~200<br \> ICC~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

ZFAT/ZNF406 Polyclonal Antibody - Protein Information

Name ZFAT

Synonyms KIAA1485, ZFAT1, ZNF406

Function


May be involved in transcriptional regulation. Overexpression causes down-regulation of a number of genes involved in the immune response. Some genes are also up-regulated (By similarity).

Cellular Location

Nucleus. Cytoplasm, cytosol

Tissue Location

Isoform 1 is strongly expressed in placenta, spleen, kidney, testis and peripheral blood leukocytes. Expressed in CD4+ and CD8+ T-cells, CD19+ B-cells and CB14+ monocytes. Isoform 3 is strongly expressed in placenta, ovary, tonsil, CD19+ B-cells and CD14+ monocytes.



Tel: 858.875.1900 Fax: 858.875.1999



ZFAT/ZNF406 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ZFAT/ZNF406 Polyclonal Antibody - Images