

MAGD4 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57181**Specification**

MAGD4 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	O96JG8
Reactivity	Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	81378

MAGD4 Polyclonal Antibody - Additional Information**Gene ID** 728239;81557**Other Names**

Melanoma-associated antigen D4, MAGE-D4 antigen, MAGE-E1 antigen, MAGED4, KIAA1859, MAGED4A, MAGEE1

Dilution

IHC-P~~N/A<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>ICC~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

MAGD4 Polyclonal Antibody - Protein Information**Name** MAGED4**Synonyms** KIAA1859, MAGED4A, MAGEE1**Function**

May enhance ubiquitin ligase activity of RING-type zinc finger-containing E3 ubiquitin-protein ligases. Proposed to act through recruitment and/or stabilization of the Ubl-conjugating enzyme (E2) at the E3:substrate complex.

Tissue Location

Expressed only in brain and ovary among normal tissues. Isoform 1 and isoform 2 are specifically expressed in glioma cells among cancer cells. Detected in some renal cell carcinoma samples.

MAGD4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MAGD4 Polyclonal Antibody - Images