

MANSC1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57204**Specification**

MANSC1 Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	O9H8J5
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46810

MANSC1 Polyclonal Antibody - Additional Information**Gene ID** 54682**Other Names**

MANSC domain-containing protein 1, Loss of heterozygosity 12 chromosomal region 3 protein, MANSC1, LOH12CR3

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

MANSC1 Polyclonal Antibody - Protein Information**Name** MANSC1**Synonyms** LOH12CR3**Cellular Location**

Membrane; Single-pass type I membrane protein

Tissue Location

Widely expressed..


MANSC1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MANSC1 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MANSC1) Polyclonal Antibody, Unconjugated (bs-18656R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample:

Large intestine(Mouse) Cell Lysate at 40 ug

Primary: Anti-MANSC1 (bs-18656R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 44 kD

Observed band size: 49 kD