

**MBIP/MAP3K12 binding inhibitory protein 1 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP57208****Specification**

---

**MBIP/MAP3K12 binding inhibitory protein 1 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	<a href="#">Q9NS73</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39281

**MBIP/MAP3K12 binding inhibitory protein 1 Polyclonal Antibody - Additional Information****Gene ID** 51562**Other Names**

MAP3K12-binding inhibitory protein 1, MAPK upstream kinase-binding inhibitory protein, MUK-binding inhibitory protein, MBIP

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**MBIP/MAP3K12 binding inhibitory protein 1 Polyclonal Antibody - Protein Information****Name** MBIP**Function**

Inhibits the MAP3K12 activity to induce the activation of the JNK/SAPK pathway. Component of the ATAC complex, a complex with histone acetyltransferase activity on histones H3 and H4.

**Cellular Location**

Nucleus. Cytoplasm. Note=Shows a cytoplasmic localization when coexpressed with MAP3K12

**Tissue Location**

Ubiquitous. High expression seen in the heart and lung

**MBIP/MAP3K12 binding inhibitory protein 1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**MBIP/MAP3K12 binding inhibitory protein 1 Polyclonal Antibody - Images**