

CRUM2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5724b**Specification**

CRUM2 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q5IJ48
Other Accession	Q80YA8 , NP_775960.4
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1258-1285

CRUM2 Antibody (C-term) - Additional Information**Gene ID** 286204**Other Names**

Protein crumbs homolog 2, Crumbs-like protein 2, CRB2

Target/Specificity

This CRUM2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1258-1285 amino acids of human CRUM2.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CRUM2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CRUM2 Antibody (C-term) - Protein Information**Name** CRB2 ([HGNC:18688](#))

Function Apical polarity protein that plays a central role during the epithelial-to-mesenchymal transition (EMT) at gastrulation, when newly specified mesodermal cells move inside the embryo

(By similarity). Acts by promoting cell ingression, the process by which cells leave the epithelial epiblast and move inside the embryo to form a new tissue layer (By similarity). The anisotropic distribution of CRB2 and MYH10/myosin-IIb at cell edges define which cells will ingress: cells with high apical CRB2 are probably extruded from the epiblast by neighboring cells with high levels of apical MYH10/myosin-IIb (By similarity). Plays a role in the maintenance of retinal neuroepithelium organization, structural integrity, adhesion, photoreceptor polarity and retinal photoreceptor layer thickness (By similarity). May play a role in determining the length of cone photoreceptor outer segments and proliferation of late-born progenitor cells (By similarity). Also required for maintenance of the apical polarity complex during development of the cortex (By similarity). Inhibits gamma-secretase- dependent cleavage of APP and secretion of amyloid-beta peptide 40 and amyloid-beta peptide 42, and thereby inhibits gamma-secretase-dependent Notch transcription (PubMed:[20299451](#)).

Cellular Location

[Isoform 1]: Apical cell membrane {ECO:0000250|UniProtKB:Q80YA8}; Single-pass type I membrane protein. Cytoplasm {ECO:0000250|UniProtKB:Q80YA8}. Cell junction {ECO:0000250|UniProtKB:Q80YA8}. Note=O-glucosylation is required for localization at the apical plasma membrane (By similarity). Distributed in a complex anisotropic pattern on apical cell edges: the level of CRB2 on a cell edge is inversely correlated with the level of MYH10/myosin-IIb (By similarity). {ECO:0000250|UniProtKB:Q80YA8}

Tissue Location

Expressed in glomeruli, podocytes of the glomerular capillary loops, and parietal glomerular epithelial cells in the kidney (at protein level) (PubMed:27942854, PubMed:29473663). Expressed in retina, fetal eye and brain (PubMed:15851977). Also expressed in kidney, RPE/choroid, and at low levels in lung, placenta, and heart (PubMed:15851977).


CRUM2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CRUM2 Antibody (C-term) - Images





CRUM2 Antibody (C-term) (Cat. #AP5724b) western blot analysis in mouse heart tissue lysates (15ug/lane). This demonstrates the CRUM2 antibody detected CRUM2 protein (arrow).



CRUM2 Antibody (C-term) (Cat. #AP5724b) immunohistochemistry analysis in formalin fixed and paraffin embedded mouse kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CRUM2 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

CRUM2 Antibody (C-term) - Background

May play a role in polarized cells morphogenesis.

CRUM2 Antibody (C-term) - References

CRUM2 Antibody (C-term) - Citations

- [Defects of CRB2 Cause Steroid-Resistant Nephrotic Syndrome.](#)