

Methionine Sulfoxide Reductase A Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57257**Specification**

Methionine Sulfoxide Reductase A Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9UJ68
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26132

Methionine Sulfoxide Reductase A Polyclonal Antibody - Additional Information**Gene ID** 4482**Other Names**

Mitochondrial peptide methionine sulfoxide reductase, 1.8.4.11, Peptide-methionine (S)-S-oxide reductase, Peptide Met(O) reductase, Protein-methionine-S-oxide reductase, PMSR, MSRA

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Methionine Sulfoxide Reductase A Polyclonal Antibody - Protein Information**Name** MSRA**Function**

Has an important function as a repair enzyme for proteins that have been inactivated by oxidation. Catalyzes the reversible oxidation-reduction of methionine sulfoxide in proteins to methionine.

Cellular Location

[Isoform 1]: Mitochondrion. [Isoform 3]: Cytoplasm. Nucleus.

Tissue Location


Ubiquitous. Highest expression in adult kidney and cerebellum, followed by liver, heart ventricles, bone marrow and hippocampus

Methionine Sulfoxide Reductase A Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Methionine Sulfoxide Reductase A Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Methionine Sulfoxide Reductase A) Polyclonal Antibody, Unconjugated (bs-18805R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Methionine Sulfoxide Reductase A) Polyclonal Antibody, Unconjugated (bs-18805R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.