

NAB1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57340**Specification**

NAB1 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q13506
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human NAB1
Epitope Specificity	401-487/487
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
SIMILARITY	Belongs to the NAB family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Transcriptional control is in part regulated by interactions between DNA-bound transcription factors, such as Egr1/NGFI-A, and coregulatory proteins, such as NAB (for NGFI-A-binding proteins). The evolutionarily conserved NAB proteins, NAB1 and NAB2, are corepressors of Egr1/NGFI-A. Both NAB1 and NAB2 contain an amino-terminal NAB-conserved domain 1 (NCB1), which is required for binding NGFI-A, and a carboxy-terminal NCD2, which is responsible for the repressor function of NAB proteins. NAB1 requires NGFI-A to gain access to DNA, indicating that NAB1 is an active repressor that works by a direct mechanism. NAB1, which is constitutively expressed, is localized exclusively in the nucleus and may play a role in controlling processes such as cell division, differentiation and apoptosis.

NAB1 Polyclonal Antibody - Additional Information**Gene ID** 4664**Other Names**

NGFI-A-binding protein 1, EGR-1-binding protein 1, Transcriptional regulatory protein p54, NAB1

Target/Specificity

Isoform Short is found in a myeloid leukemia cell line.

Dilution

IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

NAB1 Polyclonal Antibody - Protein Information

Name NAB1

Function

Acts as a transcriptional repressor for zinc finger transcription factors EGR1 and EGR2.

Cellular Location

Nucleus.

Tissue Location

Isoform Short is found in myeloid leukemia cell line KG-1

NAB1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NAB1 Polyclonal Antibody - Images