

NAP1L5 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57356

Specification

NAP1L5 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession Q96NT1

Reactivity
Host
Rat, Pig, Dog, Bovine
Rabbit

Clonality Polyclonal
Calculated MW 20 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

from human NAP1L5

Epitope Specificity 51-132/132

Isotype IgG
Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus.

SIMILARITY Belongs to the nucleosome assembly

protein (NAP) family.

Important Note This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

NAP1L5 Polyclonal Antibody - Additional Information

Gene ID 266812

Other Names

Nucleosome assembly protein 1-like 5, Down-regulated in liver malignancy, NAP1L5, DRLM

Target/Specificity

Predominantly expressed in brain.

Dilution

IHC-P~~N/A<br \><span class</pre>

="dilution_IHC-F">IHC-F~~N/A<br \><span class

="dilution_IF">IF~~1:50~200<br \> < span class = "dilution_ICC">ICC~~N/A < br \> < span class = "dilution_ICC">ICC~~N/A < span class = "dilution_ICC">ICC~~N/A < span class = "dilution_ICC">ICC~~N/A</sp

\>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH



7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

NAP1L5 Polyclonal Antibody - Protein Information

Name NAP1L5

Synonyms DRLM

Cellular Location Nucleus.


Tissue LocationPredominantly expressed in brain.

NAP1L5 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

NAP1L5 Polyclonal Antibody - Images




Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NAP1L5) Polyclonal Antibody, Unconjugated (bs-19021R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.





Paraformaldehyde-fixed, paraffin embedded (Rat colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NAP1L5) Polyclonal Antibody, Unconjugated (bs-19021R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NAP1L5) Polyclonal Antibody, Unconjugated (bs-19021R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.