

NCRNA00114 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57374**Specification**

NCRNA00114 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q6XXX2
Host	Rabbit
Clonality	Polyclonal
Calculated MW	15753

NCRNA00114 Polyclonal Antibody - Additional Information**Other Names**

Putative uncharacterized protein encoded by LINC00114, LINC00114, C21orf24, NCRNA00114

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

NCRNA00114 Polyclonal Antibody - Protein Information

Name LINC00114

Synonyms C21orf24, NCRNA00114

Tissue Location


Expressed in a range of cell lines, including B- cell lymphoma and prostate.

NCRNA00114 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NCRNA00114 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (human liver tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NCRNA00114) Polyclonal Antibody, Unconjugated (bs-19055R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NCRNA00114) Polyclonal Antibody, Unconjugated (bs-19055R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.