

NORRIN Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57378**Specification**

NORRIN Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	Q00604
Reactivity	Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	15044

NORRIN Polyclonal Antibody - Additional Information**Gene ID** 4693**Other Names**

Norrin, Norrie disease protein, X-linked exudative vitreoretinopathy 2 protein, NDP, EVR2

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

NORRIN Polyclonal Antibody - Protein Information**Name** NDP**Synonyms** EVR2**Function**

Activates the canonical Wnt signaling pathway through FZD4 and LRP5 coreceptor. Plays a central role in retinal vascularization by acting as a ligand for FZD4 that signals via stabilizing beta-catenin (CTNNB1) and activating LEF/TCF-mediated transcriptional programs. Acts in concert with TSPAN12 to activate FZD4 independently of the Wnt- dependent activation of FZD4, suggesting the existence of a Wnt- independent signaling that also promote accumulation the beta-catenin (CTNNB1). May be involved in a pathway that regulates neural cell differentiation and proliferation. Possible role in neuroectodermal cell-cell interaction.

Cellular Location

Secreted.

Tissue Location

Expressed in the outer nuclear, inner nuclear and ganglion cell layers of the retina, and in fetal and adult brain

NORRIN Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NORRIN Polyclonal Antibody - Images