

**NMES1 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP57465****Specification**

---

**NMES1 Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">O9C002</a>
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	9617

**NMES1 Polyclonal Antibody - Additional Information****Gene ID** 84419**Other Names**

Normal mucosa of esophagus-specific gene 1 protein, Protein FOAP-11, NMES1, C15orf48

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**NMES1 Polyclonal Antibody - Protein Information****Name** NMES1**Synonyms** C15orf48**Cellular Location**

Nucleus.

**Tissue Location**

Expressed mainly in stomach, placenta, small intestine and colon, as well as in normal mucosa of esophagus. Down- regulated in esophageal squamous cell carcinoma


**NMES1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **NMES1 Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (Rat intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NMES1) Polyclonal Antibody, Unconjugated (bs-19295R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.