

**PRPSAP2 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP57509**

## Specification

---

### PRPSAP2 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	<a href="#">O60256</a>
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40926

### PRPSAP2 Polyclonal Antibody - Additional Information

**Gene ID** 5636

#### Other Names

Phosphoribosyl pyrophosphate synthase-associated protein 2, PRPP synthase-associated protein 2, 41 kDa phosphoribosypyrophosphate synthetase-associated protein, PAP41, PRPSAP2

#### Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

#### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### PRPSAP2 Polyclonal Antibody - Protein Information

**Name** PRPSAP2

#### Function

Seems to play a negative regulatory role in 5-phosphoribose 1-diphosphate synthesis.

#### Tissue Location

Ubiquitous.


### PRPSAP2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## PRPSAP2 Polyclonal Antibody - Images



### Sample:


A431(Human) Cell Lysate at 40 ug

Primary: Anti-aPRPSAP2 (bs-19410R) at 1/300 dilution


Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 41 kD


Observed band size: 41 kD



Paraformaldehyde-fixed, paraffin embedded (Rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PRPSAP2) Polyclonal Antibody, Unconjugated (bs-19410R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PRPSAP2) Polyclonal Antibody, Unconjugated (bs-19410R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PRPSAP2) Polyclonal Antibody, Unconjugated (bs-19410R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.