

NUDT13 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57554**Specification**

NUDT13 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q86X67
Reactivity	Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39688

NUDT13 Polyclonal Antibody - Additional Information**Gene ID** 25961**Other Names**

NAD(P)H pyrophosphatase NUDT13, mitochondrial, 3.6.1.22, Nucleoside diphosphate-linked moiety X motif 13, Nudix motif 13, Protein KiSS-16, NUDT13 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=18827)
target="_blank">HGNC:18827)

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

NUDT13 Polyclonal Antibody - Protein Information**Name** NUDT13 ([HGNC:18827](#))**Function**

NAD(P)H pyrophosphatase that hydrolyzes NADH into NMNH and AMP, and NADPH into NMNH and 2',5'-ADP. Has a marked preference for the reduced pyridine nucleotides. Does not show activity toward NAD-capped RNAs; the NAD-cap is an atypical cap present at the 5'-end of some RNAs.

Cellular Location

Mitochondrion {ECO:0000250|UniProtKB:Q8JZU0}.

Tissue Location

Highly expressed in metastasis-suppressed chromosome 6 melanoma hybrids.
{ECO:0000269|Ref.6}

NUDT13 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NUDT13 Polyclonal Antibody - Images