

NUDT18 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57555

Specification

NUDT18 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q6ZVK8
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human NUDT18
Epitope Specificity	21-120/323
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the Nudix hydrolase family. Contains 1 nudix hydrolase domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

The protein encoded by this gene is a member of the Nudix hydrolase family. Nudix hydrolases eliminate potentially toxic nucleotide metabolites from the cell and regulate the concentrations and availability of many different nucleotide substrates, cofactors, and signaling molecules. This protein contains a Nudix hydrolase domain and hydrolyzes oxidized forms of guanosine and deoxyguanosine diphosphates. [provided by RefSeq, Sep 2012]

NUDT18 Polyclonal Antibody - Additional Information

Gene ID 79873

Other Names

8-oxo-dGDP phosphatase NUDT18, 3.6.1.58, 2-hydroxy-dADP phosphatase, 7, 8-dihydro-8-oxoguanine phosphatase, MutT homolog 3, Nucleoside diphosphate-linked moiety X motif 18, Nudix motif 18, NUDT18, MTH3

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

NUDT18 Polyclonal Antibody - Protein Information

Name NUDT18 ([HGNC:26194](#))


Function

Mediates the hydrolysis of oxidized nucleoside diphosphate derivatives. Hydrolyzes 8-oxo-7,8-dihydroguanine (8-oxo-Gua)-containing deoxyribo- and ribonucleoside diphosphates to the monophosphates. Hydrolyzes 8-oxo-dGDP and 8-oxo-GDP with the same efficiencies. Also hydrolyzes 8-OH-dADP and 2-OH-dADP. Exhibited no or minimal hydrolysis activity against 8-oxo-dGTP, 8-oxo-GTP, dGTP, GTP, dGDP and GDP. Probably removes oxidized guanine nucleotides from both the DNA and RNA precursor pools.

NUDT18 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NUDT18 Polyclonal Antibody - Images

Sample:

Testis (Mouse) Lysate at 40 ug
Primary: Anti-NUDT18 (bs-19514R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 35 kD
Observed band size: 27 kD