

PSMG2/HCCA3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57583**Specification**

PSMG2/HCCA3 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	Q969U7
Reactivity	Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29396

PSMG2/HCCA3 Polyclonal Antibody - Additional Information**Gene ID** 56984**Other Names**

Proteasome assembly chaperone 2, PAC-2, Hepatocellular carcinoma-susceptibility protein 3, Tumor necrosis factor superfamily member 5-induced protein 1, PSMG2
{ECO:0000312|EMBL:AAH13356.1}

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PSMG2/HCCA3 Polyclonal Antibody - Protein Information**Name** PSMG2 {ECO:0000312|EMBL:AAH13356.1}**Function**

Chaperone protein which promotes assembly of the 20S proteasome as part of a heterodimer with PSMG1. The PSMG1-PSMG2 heterodimer binds to the PSMA5 and PSMA7 proteasome subunits, promotes assembly of the proteasome alpha subunits into the heteroheptameric alpha ring and prevents alpha ring dimerization.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9EST4}.

Tissue Location

Widely expressed with highest levels in lung, brain and colon. Moderately expressed in muscle, stomach, spleen and heart Weakly expressed in small intestine, pancreas and liver. Highly expressed in hepatocellular carcinomas with low levels in surrounding liver tissue.

PSMG2/HCCA3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PSMG2/HCCA3 Polyclonal Antibody - Images