

### **SEC23 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57600

### **Specification**

# **SEC23 Polyclonal Antibody - Product Information**

Application IHC-P, IHC-F, IF, ICC

Primary Accession <u>Q15436</u>

Reactivity Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 86161

# **SEC23 Polyclonal Antibody - Additional Information**

### **Gene ID** 10484

#### **Other Names**

Protein transport protein Sec23A, hSec23A, SEC23-related protein A, SEC23A (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=10701" target="blank">HGNC:10701</a>)

# **Dilution**

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution\_IHC-F">IHC-F~~N/A</span> <br \> <span class
="dilution\_IF">IF~~1:50~200</span> <br \> <span class = "dilution\_ICC">ICC~~N/A</span>

# **Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

# **SEC23 Polyclonal Antibody - Protein Information**


# Name SEC23A (HGNC:10701)

### **Function**

Component of the coat protein complex II (COPII) which promotes the formation of transport vesicles from the endoplasmic reticulum (ER). The coat has two main functions, the physical deformation of the endoplasmic reticulum membrane into vesicles and the selection of cargo molecules for their transport to the Golgi complex. Required for the translocation of insulin-induced glucose transporter SLC2A4/GLUT4 to the cell membrane (By similarity).

#### **Cellular Location**

Cytoplasmic vesicle, COPII-coated vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytosol. Note=Enriched at endoplasmic reticulum exit sites, also known as transitional endoplasmic reticulum (tER)





**Tissue Location** Ubiquitously expressed.

# **SEC23 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

SEC23 Polyclonal Antibody - Images