

# SF20 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57617

## **Specification**

### SF20 Polyclonal Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

IHC-P, IHC-F, IF, ICC Q969H8 Rat, Pig, Dog

Rabbit Polyclonal 18795

# SF20 Polyclonal Antibody - Additional Information

### Gene ID 56005

#### **Other Names**

Myeloid-derived growth factor {ECO:0000303|PubMed:25581518, ECO:0000312|HGNC:HGNC:16948}, MYDGF, MYDGF (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=16948" target="\_blank">HGNC:16948</a>)

#### **Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

#### **Storage**

Store at -20  $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.

# **SF20 Polyclonal Antibody - Protein Information**

## Name MYDGF (HGNC:16948)

#### **Function**

Bone marrow-derived monocyte and paracrine-acting protein that promotes cardiac myocyte survival and adaptive angiogenesis for cardiac protection and/or repair after myocardial infarction (MI). Stimulates endothelial cell proliferation through a MAPK1/3-, STAT3- and CCND1-mediated signaling pathway. Inhibits cardiac myocyte apoptosis in a PI3K/AKT-dependent signaling pathway (By similarity). Involved in endothelial cell proliferation and angiogenesis (PubMed:<a href="http://www.uniprot.org/citations/25581518" target="blank">25581518</a>).

### **Cellular Location**

Secreted. Endoplasmic reticulum-Golgi intermediate compartment. Endoplasmic reticulum. Golgi apparatus. Note=The C-terminal RTEL motif may provide retention in the endoplasmic reticulum

### **Tissue Location**

Expressed in eosinophils (at protein level) (PubMed:29954947). Expressed in bone marrow cells



(PubMed:25581518) Expressed in synovial tissue. Found in synovial fluid of patients with arthropaties (PubMed:17362502).

# **SF20 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

SF20 Polyclonal Antibody - Images