

## SF3A3 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57619

# **Specification**

# **SF3A3 Polyclonal Antibody - Product Information**

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession Q12874

Reactivity
Host
Clonality
Calculated MW
Physical State

Rat, Pig, Dog, Bovine
Rabbit
Polyclonal
S9 KDa
Liquid

Immunogen KLH conjugated synthetic peptide derived

laG

from human SF3A3

Epitope Specificity 21-120/501

Isotype Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus speckle.

SIMILARITY

Belongs to the SF3A3 family.

SUBUNIT

Identified in the spliceosome C

JBUNIT

Identified in the spliceosome C complex.

Component of splicing factor SF3A which is composed of three subunits; SF3A3/SAP61, SF3A2/SAP62, SF3A1/SAP114. SF3A associates with the splicing factor SF3B and a 12S RNA unit to form the U2 small

and a 12S RNA unit to form the U2 small nuclear ribonucleoproteins complex (U2 snRNP). Interacts with SF3A1, through its

N-terminal region.

Important Note

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

## **Background Descriptions**

This gene encodes subunit 3 of the splicing factor 3a protein complex. The splicing factor 3a heterotrimer includes subunits 1, 2 and 3 and is necessary for the in vitro conversion of 15S U2 snRNP into an active 17S particle that performs pre-mRNA splicing. Subunit 3 interacts with subunit 1 through its amino-terminus while the zinc finger domain of subunit 3 plays a role in its binding to the 15S U2 snRNP. This gene has a pseudogene on chromosome 20. [provided by RefSeq, Jul 2008]

# SF3A3 Polyclonal Antibody - Additional Information

**Gene ID** 10946

**Other Names** 



Splicing factor 3A subunit 3, SF3a60, Spliceosome-associated protein 61, SAP 61, SF3A3, SAP61

# Target/Specificity Ubiquitous.

#### **Dilution**

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution\_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution\_IF">IF~~1:50~200</span><br \> <span class = "dilution\_ICC">ICC~~N/A</span><br \> <span class = "dilution\_E">E~~N/A</span>

#### **Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

#### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### **SF3A3 Polyclonal Antibody - Protein Information**

Name SF3A3

Synonyms SAP61

#### **Function**

Component of the 17S U2 SnRNP complex of the spliceosome, a large ribonucleoprotein complex that removes introns from transcribed pre-mRNAs (PubMed:<a

href="http://www.uniprot.org/citations/10882114" target="\_blank">10882114</a>, PubMed:<a href="http://www.uniprot.org/citations/11533230" target="\_blank">11533230</a>, PubMed:<a href="http://www.uniprot.org/citations/32494006" target="\_blank">32494006</a>, PubMed:<a href="http://www.uniprot.org/citations/34822310" target="\_blank">34822310</a>, PubMed:<a href="http://www.uniprot.org/citations/8022796" target="\_blank">8022796</a>). The 17S U2 SnRNP complex (1) directly participates in early spliceosome assembly and (2) mediates recognition of the intron branch site during pre-mRNA splicing by promoting the selection of the pre-mRNA branch-site adenosine, the nucleophile for the first step of splicing (PubMed:<a href="http://www.uniprot.org/citations/10882114" target="\_blank">10882114</a>, PubMed:<a href="http://www.uniprot.org/citations/11533230" target="\_blank">32494006</a>, PubMed:<a href="http://www.uniprot.org/citations/32494006" target="\_blank">32494006</a>, PubMed:<a href="http://www.uniprot.org/citations/34822310" target="\_blank">34822310</a>, Pu

 $href="http://www.uniprot.org/citations/10882114" target="\_blank">10882114</a>, PubMed:<a href="http://www.uniprot.org/citations/11533230" target="\_blank">11533230</a>). Involved in pre-mRNA splicing as a component of pre-catalytic spliceosome 'B' complexes (PubMed:<a href="http://www.uniprot.org/citations/29360106" target="_blank">29360106</a>, PubMed:<a href="http://www.uniprot.org/citations/30315277" target="_blank">30315277</a>).$ 

# **Cellular Location**Nucleus speckle. Nucleus

Tissue Location Ubiquitous..



Tel: 858.875.1900 Fax: 858.875.1999



# SF3A3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

SF3A3 Polyclonal Antibody - Images