

QSOX2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57627**Specification**

QSOX2 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	Q6ZRP7
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	77529

QSOX2 Polyclonal Antibody - Additional Information**Gene ID** 169714**Other Names**

Sulfhydryl oxidase 2, 1.8.3.2, Neuroblastoma-derived sulfhydryl oxidase, Quiescin Q6-like protein 1, QSOX2, QSCN6L1, SOXN

Dilution

IHC-P ~ ~ N/A
IHC-F ~ ~ N/A
IF ~ ~ 1:50 ~ 200
ICC ~ ~ N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

QSOX2 Polyclonal Antibody - Protein Information**Name** QSOX2**Synonyms** QSCN6L1, SOXN**Function**

Catalyzes the oxidation of sulfhydryl groups in peptide and protein thiols to disulfides with the reduction of oxygen to hydrogen peroxide. May contribute to disulfide bond formation in a variety of secreted proteins. Also seems to play a role in regulating the sensitization of neuroblastoma cells for interferon-gamma-induced apoptosis.

Cellular Location

Membrane; Single- pass membrane protein. Secreted. Cell membrane; Single-pass membrane protein. Nucleus membrane; Single- pass membrane protein. Note=Seems to be predominantly

targeted to the nuclear and outer plasma membrane

Tissue Location


Expressed in pancreas, brain, placenta, kidney, heart and fetal tissues. Weakly expressed in lung, liver and skeletal muscles.

QSOX2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

QSOX2 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (QSOX2) Polyclonal Antibody, Unconjugated (bs-19698R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.