

R3HDM2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57631**Specification**

R3HDM2 Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	Q9Y2K5
Host	Rabbit
Clonality	Polyclonal
Calculated MW	106999

R3HDM2 Polyclonal Antibody - Additional Information**Gene ID** 22864**Other Names**

R3H domain-containing protein 2, R3HDM2, KIAA1002

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

R3HDM2 Polyclonal Antibody - Protein Information**Name** R3HDM2**Synonyms** KIAA1002**Cellular Location**

Nucleus.


R3HDM2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)


R3HDM2 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (R3HDM2) Polyclonal Antibody, Unconjugated (bs-19704R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (R3HDM2) Polyclonal Antibody, Unconjugated (bs-19704R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

**Sample:**

A431(human) Cell Lysate at 40 ug

Raji(human) Cell Lysate at 40 ug

Primary: Anti-R3HDM2 (bs-19704R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 60,65 kD

Observed band size: 60,65 kD