

Endophilin II Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57649**Specification**

Endophilin II Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q99961
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41490

Endophilin II Polyclonal Antibody - Additional Information**Gene ID** 6455**Other Names**

Endophilin-A2, EEN fusion partner of MLL, Endophilin-2, Extra eleven-nineteen leukemia fusion gene protein, EEN, SH3 domain protein 2B, SH3 domain-containing GRB2-like protein 1, SH3GL1, CNSA1, SH3D2B

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Endophilin II Polyclonal Antibody - Protein Information**Name** SH3GL1**Synonyms** CNSA1, SH3D2B**Function**

Implicated in endocytosis. May recruit other proteins to membranes with high curvature (By similarity).

Cellular Location

Cytoplasm. Early endosome membrane; Peripheral membrane protein. Cell projection, podosome. Note=Associated with postsynaptic endosomes in hippocampal neurons.

Tissue Location

Ubiquitous. Higher expression in pancreas, placenta, prostate, testis and uterus

Endophilin II Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Endophilin II Polyclonal Antibody - Images