

### **SH3BP4 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57651

### **Specification**

## **SH3BP4 Polyclonal Antibody - Product Information**

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat, Bovine
Host
Clonality
Calculated MW
Physical State

O9P0V3
Rat, Bovine
Rabbit
Polyclonal
Liquid

Immunogen KLH conjugated synthetic peptide derived

laG

from human SH3BP4

Epitope Specificity 561-660/963

Isotype Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

SUBCELLULAR LOCATION Proclin300 and 50% Glycerol.

Membrane > clathrin-coated pit.

Cytoplasmic vesicle > clathrin-coated vesicle. Nucleus. Specifically associated with transferrin receptor-containing clathrin-coated pits and clathrin-coated vesicles. May also localize to the nucleus.

SIMILARITY Contains 1 SH3 domain.

Post-translational modifications Phosphorylated upon EGF stimulation.

Phosphorylation prevents interaction with

DNM2.

Important Note

This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

### **Background Descriptions**

This gene encodes a protein with 3 Asn-Pro-Phe (NPF) motifs, an SH3 domain, a PXXP motif, a bipartite nuclear targeting signal, and a tyrosine phosphorylation site. This protein is involved in cargo-specific control of clathrin-mediated endocytosis, specifically controlling the internalization of a specific protein receptor. [provided by RefSeq, Jul 2008]

# SH3BP4 Polyclonal Antibody - Additional Information

**Gene ID 23677** 

#### **Other Names**

SH3 domain-binding protein 4, EH-binding protein 10, Transferrin receptor-trafficking protein, SH3BP4, BOG25, EHB10, TTP



# **Target/Specificity**

Expressed in all tissues tested with higher expression in pancreas. Expressed by retinal pigment epithelial cells (at protein level).

#### **Dilution**

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution\_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution\_IF">IF~~1:50~200</span><br \> <span class ="dilution\_ICC">ICC~~N/A</span><br \> <span class ="dilution\_E">E~~N/A</span>

## **Storage**

Store at -20  $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.

# SH3BP4 Polyclonal Antibody - Protein Information

Name SH3BP4

Synonyms BOG25, EHB10, TTP

#### **Function**

May function in transferrin receptor internalization at the plasma membrane through a cargo-specific control of clathrin-mediated endocytosis. Alternatively, may act as a negative regulator of the amino acid-induced TOR signaling by inhibiting the formation of active Rag GTPase complexes. Preferentially binds inactive Rag GTPase complexes and prevents their interaction with the mTORC1 complex inhibiting its relocalization to lysosomes and its activation. Thereby, may indirectly regulate cell growth, proliferation and autophagy.

#### **Cellular Location**

Membrane, clathrin-coated pit. Cytoplasmic vesicle, clathrin-coated vesicle. Nucleus Note=Specifically associated with transferrin receptor-containing clathrin-coated pits and clathrin-coated vesicles. May also localize to the nucleus

#### **Tissue Location**

Expressed in all tissues tested with higher expression in pancreas. Expressed by retinal pigment epithelial cells (at protein level).


#### **SH3BP4 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# SH3BP4 Polyclonal Antibody - Images






Paraformaldehyde-fixed, paraffin embedded (Mouse pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SH3BP4) Polyclonal Antibody, Unconjugated (bs-19751R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SH3BP4) Polyclonal Antibody, Unconjugated (bs-19751R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat prostate); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SH3BP4) Polyclonal Antibody, Unconjugated (bs-19751R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.