

**Histone binding protein/SLBP Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP57663****Specification**

---

**Histone binding protein/SLBP Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">Q14493</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Histone binding protein/SLBP 121-220/270
Epitope Specificity	
<b>Purity</b>	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Nucleus. Polyribosome-associated. Localizes predominantly in the nucleus at the G1/G2 phases and the beginning of S phase. Through the S phase, partially redistributes to the cytoplasm. Binding to histone mRNA is necessary for cytoplasmic localization. Shuttles between the nucleus and the cytoplasm. Imported in the nucleus by the Importin alpha/Importin beta receptor. Belongs to the SLBP family.
SIMILARITY	Phosphorylated on Thr-61 and Thr-62 in the S-phase. Phosphorylation of Thr-62 by CDK1 primes phosphorylation of Thr-61 by CK2. Phosphorylation of Thr-62 is required for its degradation by the proteasome at the end of the S phase. Its degradation is not required for histone mRNA degradation at the end of the S phase. All the phosphorylated forms detected are present in the cytoplasm. Both unphosphorylated and phosphorylated forms bind the stem-loop structure of histone mRNAs.
DISEASE	Regulated during the cell cycle: protein levels increase 10 to 20 fold in the late G1 and decrease at the S/G2 border.
Important Note	This product as supplied is intended for research use only, not for use in human,

**therapeutic or diagnostic applications.****Background Descriptions**

This gene encodes a protein that binds to the stem-loop structure in replication-dependent histone mRNAs. Histone mRNAs do not contain introns or polyadenylation signals, and are processed by endonucleolytic cleavage. The stem-loop structure is essential for efficient processing but this structure also controls the transport, translation and stability of histone mRNAs. Expression of the protein is regulated during the cell cycle, increasing more than 10-fold during the latter part of G1. [provided by RefSeq, Jul 2008]

**Histone binding protein/SLBP Polyclonal Antibody - Additional Information****Gene ID** 7884**Other Names**

Histone RNA hairpin-binding protein, Histone stem-loop-binding protein, SLBP, HBP

**Target/Specificity**

Widely expressed.

**Dilution**

<span class = "dilution\_IHC-P">IHC-P~~N/A</span><br \><span class = "dilution\_IHC-F">IHC-F~~N/A</span><br \><span class = "dilution\_IF">IF~~1:50~200</span><br \><span class = "dilution\_ICC">ICC~~N/A</span><br \><span class = "dilution\_E">E~~N/A</span>

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**Histone binding protein/SLBP Polyclonal Antibody - Protein Information****Name** SLBP**Synonyms** HBP**Function**

RNA-binding protein involved in the histone pre-mRNA processing (PubMed:<a href="http://www.uniprot.org/citations/12588979" target="\_blank">12588979</a>, PubMed:<a href="http://www.uniprot.org/citations/19155325" target="\_blank">19155325</a>, PubMed:<a href="http://www.uniprot.org/citations/8957003" target="\_blank">8957003</a>, PubMed:<a href="http://www.uniprot.org/citations/9049306" target="\_blank">9049306</a>). Binds the stem-loop structure of replication-dependent histone pre-mRNAs and contributes to efficient 3'-end processing by stabilizing the complex between histone pre-mRNA and U7 small nuclear ribonucleoprotein (snRNP), via the histone downstream element (HDE) (PubMed:<a href="http://www.uniprot.org/citations/12588979" target="\_blank">12588979</a>, PubMed:<a href="http://www.uniprot.org/citations/19155325" target="\_blank">19155325</a>, PubMed:<a href="http://www.uniprot.org/citations/8957003" target="\_blank">8957003</a>, PubMed:<a href="http://www.uniprot.org/citations/9049306" target="\_blank">9049306</a>). Plays an important role in targeting mature histone mRNA from the nucleus to the cytoplasm and to the translation machinery (PubMed:<a href="http://www.uniprot.org/citations/12588979" target="\_blank">12588979</a>)

target="\_blank">12588979</a>, PubMed:<a href="http://www.uniprot.org/citations/19155325" target="\_blank">19155325</a>, PubMed:<a href="http://www.uniprot.org/citations/8957003" target="\_blank">8957003</a>, PubMed:<a href="http://www.uniprot.org/citations/9049306" target="\_blank">9049306</a>). Stabilizes mature histone mRNA and could be involved in cell-cycle regulation of histone gene expression (PubMed:<a href="http://www.uniprot.org/citations/12588979" target="\_blank">12588979</a>, PubMed:<a href="http://www.uniprot.org/citations/19155325" target="\_blank">19155325</a>, PubMed:<a href="http://www.uniprot.org/citations/8957003" target="\_blank">8957003</a>, PubMed:<a href="http://www.uniprot.org/citations/9049306" target="\_blank">9049306</a>). Involved in the mechanism by which growing oocytes accumulate histone proteins that support early embryogenesis (By similarity). Binds to the 5' side of the stem-loop structure of histone pre-mRNAs (By similarity).

### Cellular Location

Cytoplasm. Nucleus. Note=Polyribosome-associated (PubMed:12588979). Localizes predominantly in the nucleus at the G1/G2 phases and the beginning of S phase (PubMed:12588979). Through the S phase, partially redistributes to the cytoplasm (PubMed:12588979) Binding to histone mRNA is necessary for cytoplasmic localization (PubMed:12588979). Shuttles between the nucleus and the cytoplasm (PubMed:15829567). Imported in the nucleus by the Importin alpha/Importin beta receptor (PubMed:15829567)

### Tissue Location


Widely expressed..

## Histone binding protein/SLBP Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Histone binding protein/SLBP Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20

minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SLBP) Polyclonal Antibody, Unconjugated (bs-19791R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.