

Reptin/TIP49B/RUVB2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57680**Specification****Reptin/TIP49B/RUVB2 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	O9Y230
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51157

Reptin/TIP49B/RUVB2 Polyclonal Antibody - Additional Information**Gene ID** 10856**Other Names**

RuvB-like 2, 3.6.4.12, 48 kDa TATA box-binding protein-interacting protein, 48 kDa TBP-interacting protein, 51 kDa erythrocyte cytosolic protein, ECP-51, INO80 complex subunit J, Repressing pontin 52, Reptin 52, TIP49b, TIP60-associated protein 54-beta, TAP54-beta, RUVBL2, INO80J, TIP48, TIP49B

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Reptin/TIP49B/RUVB2 Polyclonal Antibody - Protein Information**Name** RUVBL2**Synonyms** INO80J, TIP48, TIP49B**Function**

Possesses single-stranded DNA-stimulated ATPase and ATP- dependent DNA helicase (5' to 3') activity; hexamerization is thought to be critical for ATP hydrolysis and adjacent subunits in the ring- like structure contribute to the ATPase activity (PubMed:10428817, PubMed:<a

[17157868](http://www.uniprot.org/citations/17157868), PubMed:<[33205750](http://www.uniprot.org/citations/33205750)>). Component of the NuA4 histone acetyltransferase complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A (PubMed:<[14966270](http://www.uniprot.org/citations/14966270)>). This modification may both alter nucleosome-DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription (PubMed:<[14966270](http://www.uniprot.org/citations/14966270)>). This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair (PubMed:<[14966270](http://www.uniprot.org/citations/14966270)>). The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400 (PubMed:<[14966270](http://www.uniprot.org/citations/14966270)>). NuA4 may also play a direct role in DNA repair when recruited to sites of DNA damage (PubMed:<[14966270](http://www.uniprot.org/citations/14966270)>). Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AZ1 from the nucleosome (PubMed:<[24463511](http://www.uniprot.org/citations/24463511)>). Proposed core component of the chromatin remodeling INO80 complex which exhibits DNA- and nucleosome-activated ATPase activity and catalyzes ATP- dependent nucleosome sliding (PubMed:<[16230350](http://www.uniprot.org/citations/16230350)>, PubMed:<[21303910](http://www.uniprot.org/citations/21303910)>). Plays an essential role in oncogenic transformation by MYC and also modulates transcriptional activation by the LEF1/TCF1-CTNNB1 complex (PubMed:<[10882073](http://www.uniprot.org/citations/10882073)>, PubMed:<[16014379](http://www.uniprot.org/citations/16014379)>). May also inhibit the transcriptional activity of ATF2 (PubMed:<[11713276](http://www.uniprot.org/citations/11713276)>). Involved in the endoplasmic reticulum (ER)-associated degradation (ERAD) pathway where it negatively regulates expression of ER stress response genes (PubMed:<[25652260](http://www.uniprot.org/citations/25652260)>). May play a role in regulating the composition of the U5 snRNP complex (PubMed:<[28561026](http://www.uniprot.org/citations/28561026)>).

Cellular Location

Nucleus matrix. Nucleus, nucleoplasm. Cytoplasm. Membrane. Dynein axonemal particle {ECO:0000250|UniProtKB:Q9DE27} Note=Mainly localized in the nucleus, associated with nuclear matrix or in the nuclear cytosol. Although it is also present in the cytoplasm and associated with the cell membranes

Tissue Location

Ubiquitously expressed. Highly expressed in testis and thymus.


Reptin/TIP49B/RUVB2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

Reptin/TIP49B/RUVB2 Polyclonal Antibody - Images



Sample: bone (mouse) Lysate at 40 ug
Primary: Anti-Reptin/TIP49B/RUVB2(bs-19836R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 51 kD
Observed band size: 51 kD



Sample:
Thymus (Mouse) Lysate at 40 ug
Primary: Anti- Reptin (bs-19836R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 51 kD
Observed band size: 54 kD