

PC5/PC6 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57708

Specification

PC5/PC6 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q92824
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	194,101 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human PC5/PC6
Epitope Specificity	361-460/1860
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
SIMILARITY	Belongs to the peptidase S8 family. Contains 1 homo B/P domain. Contains 1 PLAC domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an initial autocatalytic processing event in the ER to generate a heterodimer which exits the ER. It then sorts to the trans-Golgi network where a second autocatalytic event takes place and the catalytic activity is acquired. This encoded protein is widely expressed and one of the seven basic amino acid-specific members which cleave their substrates at single or paired basic residues. It mediates posttranslational endoproteolytic processing for several integrin alpha subunits and is thought to process prorenin, pro-membrane type-1 matrix metalloproteinase and HIV-1 glycoprotein gp160. Alternative splicing results in multiple transcript variants, some of which encode distinct isoforms, including a protease packaged into dense core granules (PC5A) and a type 1 membrane bound protease (PC5B). [provided by RefSeq, May 2014]

PC5/PC6 Polyclonal Antibody - Additional Information

Gene ID 5125

Other Names

Proprotein convertase subtilisin/kexin type 5, 3.4.21.-, Proprotein convertase 5, PC5, Proprotein

convertase 6, PC6, hPC6, Subtilisin/kexin-like protease PC5, PCSK5, PC5, PC6

Target/Specificity

Expressed in T-lymphocytes.

Dilution

IHC-P~~N/A
IHC-F~~N/A
IF~~1:50~200
ICC~~N/A
E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PC5/PC6 Polyclonal Antibody - Protein Information

Name PCSK5

Synonyms PC5, PC6

Function

Serine endoprotease that processes various proproteins by cleavage at paired basic amino acids, recognizing the RXXX[KR]R consensus motif. Likely functions in the constitutive and regulated secretory pathways. Plays an essential role in pregnancy establishment by proteolytic activation of a number of important factors such as BMP2, CALD1 and alpha-integrins.

Cellular Location

[Isoform PC6A]: Secreted. Note=Secreted through the regulated secretory pathway.

Tissue Location

Expressed in T-lymphocytes.

PC5/PC6 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PC5/PC6 Polyclonal Antibody - Images